

such as the production of nutritional supplements and the conversion of carbon to chemical intermediates such as ethanol. Ethanol can be used to make synthetic aviation fuel (SAF) and other valuable chemicals.

Enzymes for energy efficient solvent-based CO, capture Hot potassium carbonate (HPC) applied to catalyse CO, absorption in the HPC process through the use of carbonic anhydrase (CA). This is an enzyme which plays an essential role in humans and many animals. Its function is to remove CO₂, which is a by-product of respiration.

Respiration is the mechanism by which animals convert oxygen that they breathe and food that they eat to energy. systems. The molecular structure

supports that by catalysing the reaction of CO, with water to form bicarbonate and hydrogen ions:

 $CO_2 + H_2O \rightarrow HCO3-+ H+.$

A proprietary CA enzyme known as 1T1 is used by CO, Solutions by Saipem on industrial CO, capture

potassium carbonate solution and whilst also allowing low-cost manufacture of the enzyme.

Algae for CO, capture

Algae use photosynthesis to grow their cell mass, however unlike trees and plants, their entire function is dedicated to photosynthesis and they do not expend energy growing

physical structures. The implication is that per acre of land, they are much more efficient at capturing CO, than

The British start-up Brilliant Planet has also developed a CO, capture process that relies on algae. However, its intent is not to utilise the products

location. Seawater is pumped into a lagoon filled with algae. Nutrients in the seawater are absorbed by the algae as they grow. They simultaneously absorb CO, from the air in a form of direct air capture (DAC) of CO₂.

CO, capture is also achieved through a second mechanism. The algae consume bicarbonate ions from the seawater. These ions have been

28 | gasworld • February 2024 gasworld.com February 2024 • gasworld | 29



▶ created naturally during absorption of atmospheric CO₂. As the lagoon water is released back into the sea, it re-absorbs CO₂ from the atmosphere. This is also a natural form of direct air capture of CO₂.

"...a new generation of algae technologists are turning to CO₂ capture as their business model" The algae are harvested, dried and buried under sand in the desert. The dry, salty algae are acidic and cannot biodegrade in the desert sands. This guarantees that the captured CO_2 is permanently sequestered.

Omega-3 oils from algae for CO₂ capture and utilisation

Various types of algae are used in cosmetics and as a meat substitute in vegan and vegetarian foods. The use of pure CO₂ gas to accelerate the growth of algae in these commercial utilisation applications has been

common for several years. However, a new generation of algae technologists are turning to CO₂ capture as their business model.

The Danish start-up ALGIECEL has targeted the capture of biogenic CO_2 as biogas is upgraded to biomethane from anaerobic biogas reactors. Its vision is to utilise the captured CO_2 to cultivate algae which will be rich in omega-3 oils. These can be used as nutritional supplements for humans, livestock and farmed fish.

The ALGIECEL PhotoBioReactor (PBR) is a containerised unit designed •

30 | gasworld • February 2024

gasworld.com



▶ to match the scale of biogas production on many of Europe's 18,000 biogas reactors. Light for the photosynthesis is provided by energy efficient LEDs to ensure 24-hour operation at maximum CO₂ capture and conversion yields.

Bio-fermentation to ethanol

Like algae, bacteria can also consume CO₂ and carbon monoxide (CO) to produce valuable chemicals such as ethanol.

LanzaTech utilises anaerobic acetobacter bacteria in a fermenter to convert CO rich feed gases, such as syngas, to ethanol and a range of biochemicals.

The ideal feedstocks for the LanzaTech fermenter are CO-rich. Iron and steelmaking yields CO-rich flue gases which are ideal feedstocks to the LanzaTech process. Blast furnace gas (BFG) contains 20% CO and converter gas (also known as basic oxygen furnace gas or BOFG) contains 60% CO.

The energy-rich converter gas (LHV 3 kWh/Nm³) is often utilised on the iron and steelmaking facility for heat or power generation on a gas engine. Alternatively, it is often flared. Blast furnace gas has a lower energy value due to the lean CO concentration and higher CO₂ content (LHV 0.9 kWh/Nm³). It can also be always utilised on the facility or is sent to the flare. Utilisation of the BFG in the LanzaTech process can generate valuable bioethanol.

LanzaTech's process was demonstrated at pilot-scale in 2008 using flue gases from the BlueScope Steel mill in Glenbrook, New Zealand. Since then, LanzaTech has successfully deployed its technology at two 300 tonnes per annum demonstration facilities at Baosteel Shanghai and Shougang Steel Caofeidian in China. These LanzaTech fermenters are fed with a range of iron and steelmaking off-gases including BOFG, BFG, and coke oven gas (COG).


Ethanol as a fuel and SAF precursor

The term CCT or 'Carbon Capture and Transformation' has been used to describe the LanzaTech fermentation process. It transforms captured carbon to ethanol. Subsequently, the complimentary LanzaJet process can

be used to convert the bioethanol to synthetic aviation fuel (SAF) in its proprietary 'Ethanol to Jet' or ETJ process.

LanzaTech operates the Freedom
Pines Biorefinery in Soperton, Georgia
which uses bio-fermenters to generate
ethanol and other chemicals. The
LanzaJet ethanol to jet SAF production
process will soon also be implemented
at that location to utilise captured
carbon to make fuels that can substitute
aviation kerosene, a fossil fuel distilled
from crude oil.

Ethanol is also a fuel component in its own right. In Europe it is blended with gasoline at either 5 to 10%. In the US, the ethanol composition can be up to 15%. In China, there is also a gasoline type fuel that is 85% ethanol with a 15% gasoline fraction.



32 | gasworld • February 2024 gasworld.com